

Bivalve aquaculture in Croatia—historical perspective


Melita Peharda & Daria Ezgeta-Balić

Institute of Oceanography and Fisheries (Split, Croatia)

www.izor.hr



Tradition
Mali Ston Bay



- In early 1940s increased production—problem with sales, very limited export
- WWII—all facilities devastated
- 1946. Fisheries Station—science and technology development




Source: Basaric 1981. Logor kraljevske na morskoj obali Jadranskog mora, s posebnim osvrtom na Malostonski zaliv. "Malostonski zaljev prirađen podlogu i društveno valoriziranje." JAZU - Dnevnik, pp. 268-281.



Tradition
Mali Ston Bay



Mediterranean mussel
(Mytilus galloprovincialis)
Commercial size > 5 cm
Production cycle 1.5-3 years








European flat oyster
(Ostrea edulis)
Commercial size > 6 cm
Production cycle 2-3 years

Two species ... Two stories




From north to south...

Lim channel...Novigrad Sea...Krka River Estuary
Mali Ston Bay



Annual production (1000 tons)




Annual production (1000 tons)

Long term trends...

Europe vs. Croatia; Oyster vs. Mussel

STRENGTHS	WEAKNESSES
<ul style="list-style-type: none"> Pristine environment <i>Ostrea edulis</i> populations High market demands 	<ul style="list-style-type: none"> Demography of producers Lack of hatchery Administration requirements
OPPORTUNITIES	THREATS
<ul style="list-style-type: none"> Multitrophic aquaculture Branding of products Increasing demand (tourism) 	<ul style="list-style-type: none"> Diseases Coastal development Invasive species (<i>Crassostrea gigas</i>)



— France
— Spain
— Ireland
— Croatia
— Greece

